


---

Homework 1  
Traitement Quantique de l'Information

---

**Exercise 1** *The Young double slit experiment (1803)*

In this exercise we want to calculate the form of the Young interference fringes. A beam of monochromatic light of wave length  $\lambda$  is sent through a double slit, and the light is reflected on a screen at a distance  $D$ . The distance between the two slits is  $d$ .



We assume that the waves diffracted by each slit have a spherical shape ( $\lambda$  the wavelength and  $\nu$  the frequency):

$$\phi_B(\vec{r}) = A \frac{e^{i\left(\frac{2\pi}{\lambda}|\vec{r}_B - \vec{r}\right] - 2\pi\nu t}}{|\vec{r}_B - \vec{r}|}, \quad \phi_C(\vec{r}) = A \frac{e^{i\left(\frac{2\pi}{\lambda}|\vec{r}_C - \vec{r}\right] - 2\pi\nu t}}{|\vec{r}_C - \vec{r}|}.$$

The total wave function at  $P$  on the screen is

$$\psi(\vec{r}_P) = \phi_B(\vec{r}_P) + \phi_C(\vec{r}_P).$$

We will use the plane wave approximation for  $D \gg d$ :

$$\psi(\vec{r}_P) \simeq \frac{A}{D} e^{-2\pi i \nu t} \left( e^{\frac{2\pi i}{\lambda} |\vec{r}_B - \vec{r}_P|} + e^{\frac{2\pi i}{\lambda} |\vec{r}_C - \vec{r}_P|} \right).$$

1) Show that the intensity at  $P$  on the screen is equal to

$$|\psi(\vec{r}_P)|^2 \approx \frac{4A^2}{D^2} \cos^2 \left( \frac{\pi d}{\lambda} \sin \theta \right).$$

Hint: evaluate first the path difference  $|\vec{r}_C - \vec{r}_P| - |\vec{r}_B - \vec{r}_P|$  for  $D \gg d$ .

2) Find the condition on  $\sin \theta$  which leads to minima and maxima of the intensity on the screen.

- 3) Let  $\rho$  be the coordinate on the screen measured from  $O$ . We have  $\tan \theta \approx \frac{\rho}{D}$  and since  $\theta$  is small  $\theta \approx \frac{\rho}{D}$ . Compute the distance between two successive minima of the intensity pattern on the screen.  
Let  $d = 0.25\text{mm}$ ,  $D = 10\text{m}$  and  $\lambda = 652\text{nm}$  (red light). What is the distance between two successive minima?

### Exercise 2 *Modern Young's experiment*

Young's double slit experiment has been reformed with Carbon 60 molecules,  $C_{60}$  in 1999. Surprisingly, these molecules behave like waves when they are well isolated from their environment. The more recent experiments have evidenced such a wave-like behavior for bigger molecules with 400 to 1000 atoms.

The diameter of  $C_{60}$  (this molecule has the form of a sphere and contains 60 carbon atoms) is approximately 0.7 nm and a mole containing  $N_A = 6.022 \times 10^{23}$  carbon atoms weights 12 grams.

- 1) Compute the De Broglie wavelength of molecules produced in an oven which have an average velocity of 220m/s. Compare with the size of individual molecules.
- 2) We perform a Young's experiment with  $d = 100\text{nm}$  and  $D = 1.25\text{m}$ . What do we observe on the screen assuming a wave like behavior?
- 3) A football weights approximately 450g and the initial velocity of a professional shoot can attain 100km/h. Estimate the De Broglie wavelength.

### Exercise 3 *Photoelectric effect*

The maximal wavelength to extract a photoelectron from tungsten is 230nm (ultraviolet). What is the necessary wavelength of light to extract electrons with kinetic energy 1.5eV? What is the speed of these electrons?

#### Useful Constant

$$c = 2.997 \times 10^8 \text{ m/s (speed of light)}$$

$$\hbar = 1.054 \times 10^{-34} \text{ J} \cdot \text{s} \left( \frac{h}{2\pi} \text{ where } h \text{ is Planck's constant} \right)$$

$$m = 9.109 \times 10^{-31} \text{ kg (mass of an electron)}$$

$$1\text{eV} = 1.6 \times 10^{-19} \text{ J}$$