

# Nuclear Fusion and Plasma Physics - Exercises

Prof. A. Fasoli - Swiss Plasma Center / EPFL


Problem Set 1 - 09 September 2024

## Exercise 1 - Combustion, Fission or Fusion?


A Swiss household consumes 18780 kWh per year on average. With 1 kg of *fuel*, for how many years can you supply energy for one household?

Consider three cases:


- a) The conventional **chemical reaction**, which produces  $6.5 \times 10^{-19}$  J of energy per reaction.



- b) The **nuclear fission** reaction



- c) The **nuclear fusion** reaction



*Hint:* For points b) and c) use Einstein's relation:

$$E = \Delta m_0 c^2 \tag{1}$$

where  $\Delta m_0$  is the *mass deficit*, i.e. the difference between the rest masses before and after the reaction.


*Indications:*  $m_{\text{C}} = 12.011 \text{ u}$ ,  $m_{\text{O}} = 15.999 \text{ u}$ ,  $m_{\text{n}} = 1.0087 \text{ u}$ ,  $m_{\text{U}} = 235.04 \text{ u}$ ,  $m_{\text{Ce}} = 139.91 \text{ u}$ ,  $m_{\text{Zr}} = 93.91 \text{ u}$ ,  $m_{\text{D}} = 2.014 \text{ u}$ ,  $m_{\text{T}} = 3.0164 \text{ u}$ ,  $m_{\text{He}} = 4.0027 \text{ u}$ , with  $1 \text{ u} = 1.66 \cdot 10^{-27} \text{ kg}$ .

## Exercise 2 - A “small” Tokamak



Consider a Tokamak plasma uniformly distributed in a volume  $V = 10 \text{ m}^3$  with the following parameters:  $B = 2 \text{ T}$ ;  $n_e = 10^{20} \text{ m}^{-3}$ ;  $T_i = T_e = 10 \text{ keV}$ . The total heating power delivered to the plasma is  $P_{in} = 28 \text{ MW}$  and the tokamak is assumed to be in “steady state”.

a) Using the cross-sections in the figure below, calculate:

- The power in the  $\alpha$ -particle population and the total fusion power in D-T (50:50 mixture).
- The total fusion power in D-D.



*Hint:* The fusion reactions and energies of the products are:



- b) Calculate the *physics fusion gain* “ $Q$ ” and the Lawson parameter  $n_e \tau_E$ . How close are we to break-even?
- c) Making simple assumptions on the tokamak geometry (torus with major radius  $R_0 = 1 \text{ m}$  and circular cross-section), consider the situation of a D-T mixture plasma with the heat flux concentrated only in 1/10 of the wall surface. Is it suitable to cover that portion of surface with a material having a heat tolerance of  $4 \text{ MW/m}^2$ ?

## Exercise 3 - The effect of impurity contamination on the plasma power balance

1. Assuming a  $T = 10$  keV plasma with  $n_e = 2 \times 10^{20} \text{ m}^{-3}$  and a volume  $V = 10 \text{ m}^3$ , calculate the radiated power due to impurity contamination and the fractional reduction of the fusion power due to dilution of the fusion fuel, in two cases:
  - a) Carbon impurities:
    - Assume that the carbon atoms are fully ionized ( $Z = 6$ ), and  $n_{C^{6+}}/n_e = 4\%$ .
    - The radiated power function in  $T = 10$  keV, assuming coronal equilibrium, is  $R_C = 10^{-34} \text{ Wm}^3$ .
  - b) Tungsten impurities:
    - Assume that the charge state of the tungsten atoms is 50 ( $Z = 50$ ), and  $n_{W^{50+}}/n_e = 10^{-5}$ .
    - The radiated power function in  $T = 10$  keV, assuming coronal equilibrium, is  $R_W = 10^{-31} \text{ Wm}^3$ .

*Note:*

- The impurity radiation losses are calculated as  $\frac{P_{rad}^Z}{V} = n_Z n_e R_Z [W/m^3]$ , where  $R_Z$  is the radiative power function for the impurity species.
  - The plasma retains quasi-neutrality:  $n_e = n_{DT} + Z n_Z$ , where  $Z$  is the charge state of the impurity species, and  $n_Z$  the impurity density.
  - The fractional reduction of fusion power is defined as  $F = \frac{P_{norm} - P_{dilu}}{P_{norm}}$ , where  $P_{dilu}$  is the fusion power density with diluted fuel and  $P_{norm}$  is the fusion power density without dilution.
2. If we define the parameter  $\rho^*$  as the ratio between the global alpha particle confinement time and the energy confinement time, can you qualitatively justify the importance of  $\rho^*$  for plasma ignition, evident in the figure below?




Figure 1: Ignition condition in the presence of fusion-produced  $\alpha$ -particles, with varying confinement times [D. Reiter, G.H. Wolf, H. Kever, Nucl. Fusion, 30, (1990), 2141].