

# Nuclear Fusion and Plasma Physics - Exercises

Prof. A. Fasoli - Swiss Plasma Center / EPFL

Solutions to Problem Set 3 - 09 October 2023

## Exercise 1 - Plasma Production

a) The definition of **relative degree of ionization** is

$$\alpha = \frac{n_e}{n_e + n_{Ar}}$$

where  $n_{Ar} = N_{Ar}/V$  is the density of neutral Argon atoms (number of Ar atoms per  $\text{m}^3$ ). To evaluate  $n_{Ar}$  we can use the ideal gas law:

$$p_{Ar} = n_{Ar} k_B T_{Ar} \quad (1)$$

where  $p_{Ar}$  is the pressure of Argon inside the vacuum chamber in Pascal,  $k_B = 1.38 \times 10^{-23} \text{ J/K}$  is the Boltzmann constant and  $T_{Ar}$  is the temperature of the Argon gas in Kelvin (normally assumed to be at room temperature - 298 K). Inverting this equation for  $n_{Ar}$  one finds that

$$n_{Ar} = \frac{p_{Ar}}{k_B T_{Ar}} \quad (2)$$

In order to use this expression one needs to convert the pressure given in Torr to Pascal using  $760 \text{ Torr} = 1.01 \times 10^5 \text{ Pa}$ . Then  $p_{Ar} = 1.33 \times 10^{-2} \text{ Pa}$ . Thus, the neutral Argon number density  $n_{Ar}$  is

$$n_{Ar} = 3.23 \times 10^{18} \text{ m}^{-3} \quad (3)$$

The degree of ionization with  $n_e = 1 \times 10^{16} \text{ m}^{-3}$  and  $n_{Ar} = 3.23 \times 10^{18} \text{ m}^{-3}$  is

$$\alpha = \frac{n_e}{\underbrace{n_e + n_{Ar}}_{\approx n_{Ar}}} = \frac{1 \times 10^{16}}{1 \times 10^{16} + 3.23 \times 10^{18}} \approx 3.08 \times 10^{-3}$$

b) The electron-neutral **collision frequency** is

$$\nu_{en} = n_{Ar} \sigma_n v_{rel}$$

where  $v_{rel}$  is the relative velocity between electrons and neutrals and  $\sigma_n = 10^3 \pi a_0^2$  is the collision cross-section. Since  $m_e \ll m_{Ar}$  and  $T_e \gg T_0$ , we can assume  $v_{rel} \simeq v_e$ .

In general,  $\nu_{en}$  is a function of the electron velocity and, implicitly,  $\sigma_n = \sigma_n(v_e)$ . In our problem we can consider  $\sigma_n$  constant and a typical velocity of the electrons equal to their thermal velocity  $v_{the} = \sqrt{\frac{eT_e}{m_e}}$ .

Plugging these numbers in the expression above we find:

$$\nu_{en} = 3.23 \times 10^{18} \text{ m}^{-3} \cdot 10^3 \pi \underbrace{(5.29 \times 10^{-11})^2}_{a_0^2} \text{ m}^2 \cdot \sqrt{\frac{eT_e}{m_e}} \frac{\text{m}}{\text{s}} \approx 2.06 \times 10^7 \text{ s}^{-1}$$

c) Can we consider this gas to be a **plasma**?

- The Debye length is:

$$\lambda_D = \sqrt{\frac{\varepsilon_0 T_e}{e^2 n_e}} \approx 7430 \sqrt{\frac{T_e [\text{eV}]}{n_e [\text{m}^{-3}]}} = 7430 \sqrt{\frac{3}{10^{16}}} \text{ m} = 0.13 \text{ mm}$$

The ionized gas is confined in a container of dimension  $L_p \approx 0.5 \text{ m} \gg 0.13 \text{ mm}$ . We see then that  $L_p \gg \lambda_D$  as required for a plasma.

- $N_D = \frac{4}{3} \pi \lambda_D^3 n_e \approx 9.2 \times 10^4 \gg 1$ , so the condition of the plasma parameter  $g = N_D^{-1} \ll 1$  is verified.
- To see dynamic collective effects in a plasma (oscillations at the frequency  $\omega_p$ ), we need  $\omega_p$  to be much larger than the collision frequency:

$$\omega_p = \sqrt{\frac{e^2 n_e}{m_e \varepsilon_0}} \approx 18\pi \sqrt{n_e [\text{m}^{-3}]} \text{ rad/s} = 18\pi \sqrt{10^{16}} \text{ rad/s} = 5.7 \times 10^9 \text{ rad/s}$$

To compare  $\omega_p$  with  $\nu_{en}$  we need to convert it in  $\text{s}^{-1}$ :

$$f_p = \frac{\omega_p}{2\pi} \approx 0.9 \times 10^9 \text{ s}^{-1} > \nu_{en} = 2.06 \times 10^7 \text{ s}^{-1}$$

We can therefore conclude that this ionized gas *is* a plasma.

## Exercise 2 - Mirror effect

- a) In an ideal solenoid the field lines would be straight through the ring-shaped conductors. However due to the finite distance between the two coils the field lines will diverge after the first ring, and reconverge when entering the second ring.
- b) The field will be strongest at the location of the coils, and weaker in the middle. Far to the left and right of the coils the field will decay approximately as  $1/r^3$ , similarly as for a magnetic dipole.




Figure 1: Field lines and field strength in a magnetic mirror.

- c) A particle with velocity only along the axis will feel no force because it moves parallel to the magnetic field. It will continue undisturbed along its path.
- d) Denote the minimum magnetic field strength halfway the coils as  $B_0$  and the maximum field at each coil as  $B_1$ . Conservation of kinetic energy gives

$$\frac{1}{2}mv_{\parallel,0}^2 + \frac{1}{2}mv_{\perp,0}^2 = \frac{1}{2}mv_{\parallel,1}^2 + \frac{1}{2}mv_{\perp,1}^2$$

Conservation of the adiabatic invariant gives

$$\frac{mv_{\perp,0}^2}{B_0} = \frac{mv_{\perp,1}^2}{B_1}$$

Using this result, one can substitute for  $v_{\perp,1}^2$

$$\frac{1}{2}mv_{\parallel,0}^2 + \frac{1}{2}mv_{\perp,0}^2 \left(1 - \frac{B_1}{B_0}\right) = \frac{1}{2}mv_{\parallel,1}^2$$

Since  $B_1 > B_0$  the second term on the left hand side is negative. If  $B_1$  is large enough then this term will cancel the first term, and  $v_{\parallel,1}$  can become zero.




Figure 2: Loss cone of a particle in a magnetic mirror.

- e) If a particle is reflected, there must be a point on its trajectory where  $v_{\parallel,1} = 0$ . Rearranging the expression found in (d) we get

$$\frac{1}{2}mv_{\parallel,0}^2 + \frac{1}{2}mv_{\perp,0}^2 = \frac{B_1}{B_0} \frac{1}{2}mv_{\perp,0}^2 \quad (4)$$

$$\frac{v_{\parallel,0}^2 + v_{\perp,0}^2}{v_{\perp,0}^2} = \frac{B_1}{B_0} \quad (5)$$

$$\sin^2 \theta_c = \frac{B_0}{B_1} \quad (6)$$

Here  $\theta_c$  is an angle on the  $(v_{\parallel,0}, v_{\perp,0})$  plane (see Fig. 2). The zone  $\theta < \theta_c$  is referred to as the *loss cone* of the velocity distribution. Particles with velocity components in this loss cone at the midplane will not be trapped in the magnetic mirror and will escape.

The loss cone is one of the fundamental reasons why fusion based on magnetic mirrors is difficult to achieve. For confinement, we would like particles to have a high perpendicular velocity. However, due to collisions, the particles will acquire a parallel velocity as well, pushing them into the loss cone and allowing them to escape from the mirror.

## Exercise 3 - Confinement by a toroidal field

a) Ampère's law in integral form reads

$$\oint_C \mathbf{B} \cdot d\boldsymbol{\ell} = \mu_0 \iint_S \mathbf{j} \cdot d\mathbf{S} = \mu_0 I$$

This is valid for any contour containing the current carrying wire. Due to symmetry, if we choose the contour as a circle centered at the wire,  $B$  is constant along the integration path. Therefore

$$B \oint_C d\ell = 2\pi r B_\theta = \mu_0 I$$
$$B_\theta = \frac{\mu_0 I}{2\pi r}$$

Clearly the field strength decreases as  $1/r$ .

Using Ampère's law in differential form (in cylindrical coordinates) we get a similar result

$$(\nabla \times B)_z = \frac{1}{r} \left[ \frac{\partial}{\partial r}(rB_\theta) - \frac{\partial B_r}{\partial \theta} \right] = 0$$

We immediately remove terms involving  $\partial/\partial\theta$  because of symmetry.

$$\frac{1}{r} \frac{\partial}{\partial r}(rB_\theta) = 0$$
$$rB_\theta = k$$
$$B_\theta = \frac{k}{r}$$

with  $k$  the appropriate constant of integration. In this case also we see that the field strength decreases as  $1/r$ .

The gradient of the magnetic field strength is easy to compute, since the only component is  $B_\theta$  which depends only on  $r$ .

$$\nabla B = \frac{d}{dr} B_\theta(r) \hat{\mathbf{r}}$$
$$= \frac{d}{dr} \left( \frac{\mu_0 I}{2\pi r} \right) \hat{\mathbf{r}}$$
$$= -\frac{\mu_0 I}{2\pi r^2} \hat{\mathbf{r}}$$

As could be expected, the gradient is oriented in the  $-\hat{\mathbf{r}}$  direction.




Figure 3: Toroidal  $B$  field created by current carrying wire. The  $\nabla B$ -drift (which is opposite for electrons and ions) causes charge separation which creates an electric field. The resulting  $E \times B$  drift drives the bulk of the plasma outwards.

- b) We can now evaluate the direction of the various drifts. The  $\nabla B$  drift is in the direction of  $\mp B \times \nabla B$  for electrons and ions respectively. The curvature drift has direction  $\mp R_c \times B$  for electrons and ions respectively. Since  $\nabla B$  and  $R_c$  are opposite, both drifts will have the same effect: they will cause ions to drift upwards and electrons to drift downwards.

This charge separation will result in an electric field in the  $-\hat{z}$  direction:  $\mathbf{E} = -E \hat{z}$ . The resulting  $E \times B$  drift is oriented in the outward radial direction for both ions and electrons. This is the fundamental reason why it is not possible to confine a plasma in a simple toroidal field.

*Note:* in a Tokamak, this problem is solved by driving a current through the plasma. The resulting poloidal magnetic field will add to the toroidal field, producing helical twisted field lines which periodically visit both top and bottom parts of the plasma. This provides a path for the particles to counteract the charge separation, thus “short circuiting” this instability.