

**Homework 4****Exercise 1.\***

a) Let  $X$  be a *continuous* and *non-negative* random variable defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . Show that

$$\mathbb{E}(X) = \int_0^{+\infty} (1 - F_X(t)) dt.$$

Extend this formula to further to any continuous random variable  $X$ . That is, show that

$$\mathbb{E}(X) = \int_0^{+\infty} (1 - F_X(t)) dt - \int_{-\infty}^0 F_X(t) dt.$$

b) Use this formula to compute  $\mathbb{E}(X)$  for  $X \sim \text{Laplace}(0, \lambda^{-1})$  for  $\lambda > 0$ .

c) Let  $X$  be a *discrete* and *non-negative* random variable taking values in  $\mathbb{N}$  only. Show that

$$\mathbb{E}(X) = \sum_{k \geq 0} (1 - F_X(k)).$$

and use this new formula to compute  $\mathbb{E}(X)$  when  $X \sim \text{Geom}(p)$  for some  $0 < p < 1$ .

**Exercise 2.** Let  $\lambda > 0$  and  $X \sim \mathcal{E}(\lambda)$ , and let us define  $Y = X^a$ , where  $a \in \mathbb{R}$ .

a) For what values of  $a \in \mathbb{R}$  does it hold that  $\mathbb{E}(Y) < +\infty$ ?

b) For what values of  $a \in \mathbb{R}$  does it hold that  $\mathbb{E}(Y^2) < +\infty$ ?

c) For what values of  $a \in \mathbb{R}$  is  $\text{Var}(Y)$ :

c1) well-defined and finite?      c2) well-defined but infinite?      c3) ill-defined?

d) Compute  $\mathbb{E}(Y)$  and  $\text{Var}(Y)$  for the values of  $a \in \mathbb{Z}$  such that these quantities are well-defined.

*Hint:* Use integration by parts, recursively.

**Exercise 3.** Let  $X$  be a random variable that is symmetrically distributed (i.e.  $X \sim -X$ ) and square-integrable with  $\text{Var}(X) = 1$ . Let also  $Y = 1_{\{X \geq 0\}}$ .

a) Show that for any distribution of the random variable  $X$ ,  $\text{Cov}(X, Y) \geq 0$ .

b) Using the inequality  $\text{Cov}(X, Y) \leq \sqrt{\text{Var}(X)} \sqrt{\text{Var}(Y)}$  (whose proof is to come in the sequel of the course), find the least value  $C > 0$  such that  $\text{Cov}(X, Y) \leq C$  for every distribution of  $X$ .

c) Compute  $\text{Cov}(X, Y)$  for  $X \sim \mathcal{N}(0, 1)$ .

d) Is it possible to find a distribution for  $X$  such that  $\text{Cov}(X, Y) = C$ ? If not, is it possible to find a sequence of random variables  $(X_n, n \geq 1)$  with varying distributions (all respecting the above constraints) and  $Y_n = 1_{\{X_n \geq 0\}}$ , such that  $\text{Cov}(X_n, Y_n) \xrightarrow[n \rightarrow \infty]{} C$ ?

e) Is it possible to find a distribution for  $X$  such that  $\text{Cov}(X, Y) = 0$ ? If not, is it possible to find a sequence of random variables  $(X_n, n \geq 1)$  with varying distributions (all respecting the above constraints) and  $Y_n = 1_{\{X_n \geq 0\}}$ , such that  $\text{Cov}(X_n, Y_n) \xrightarrow[n \rightarrow \infty]{} 0$ ?

**Exercise 4.** a) Let  $X$  be a Poisson random variable with parameter  $\lambda > 0$ . Compute its characteristic function  $\phi_X$ .

b) Show that for a discrete random variable  $X$  with values in  $\mathbb{Z}$ , the following inversion formula holds:

$$\mathbb{P}(\{X = k\}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-itk} \phi_X(t) dt, \quad \forall k \in \mathbb{Z}$$

c) Use the above formula to deduce the distribution of the random variable  $X$  with values in  $\mathbb{Z}$  whose characteristic function is given by

$$\phi_X(t) = \cos(t), \quad t \in \mathbb{R}$$

d) Without solving part c), how could you be sure that  $\phi_X$  is indeed a characteristic function?