

Astrophysics IV : Stellar and galactic dynamics

Exercises**Problem 1 :**

In a first part, estimate the relaxation time of an ultra-faint dwarf galaxy (UFD) containing only about 1'000 stars and being as compact as 50 pc. In a second part, assume that the typical velocity of the stars is about 4 km/s and thus, that the stellar component must be embedded in a massive dark matter halo. How is the relaxation time changed from the first to the second case ?

Hint : Assume that the mass of the stars is about one solar mass.

Problem 2 :

Demonstrate that the Poisson equation can be derived from a variational principle and interpret the meaning of the extremalisation performed.

Problem 3 :

In practice, for spherical systems, is often useful to derive, for example the density $\rho(r)$ knowing the gravitational field $g(r) = -\frac{d\Phi}{dr}$, or the potential $\Phi(r)$ knowing the cumulative mass $M(r)$. Using the relations presented during the lectures, express successively $\rho(r)$, $\Phi(r)$, $M(r)$ and $\frac{d\Phi}{dr}$ as a function of respectively $\rho(r)$, $\Phi(r)$, $M(r)$ and $\frac{d\Phi}{dr}$ as given in the following table :

	$\rho(r)$	$\Phi(r)$	$M(r)$	$\frac{d\Phi}{dr}$
$\rho(r)$	$\rho(r)$	$\frac{1}{4\pi G} \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\Phi}{dr} \right)$	$\frac{1}{4\pi r^2} \frac{dM(r)}{dr}$	$\frac{1}{4\pi G} \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\Phi}{dr} \right)$
$\Phi(r)$	$-\frac{GM(r)}{r} - 4\pi G \int_r^\infty dr' r' \rho(r')$	$\Phi(r)$	$-G \int_r^\infty dr' \frac{M(r')}{r'^2}$	$-\int_r^\infty dr' \frac{d\Phi}{dr}$
$M(r)$	$4\pi \int_0^r dr' r'^2 \rho(r')$	$\frac{r^2}{G} \frac{d\Phi}{dr}$	$M(r)$	$\frac{r^2}{G} \frac{d\Phi}{dr}$
$\frac{d\Phi}{dr}$	$\frac{4\pi G}{r^2} \int_0^r dr' r'^2 \rho(r')$	$\frac{d\Phi}{dr}$	$\frac{GM(r)}{r^2}$	$\frac{d\Phi}{dr}$