

Exercise 1 *Grover's algorithm for $N = 4$*

- (a) The theory says that one query of the oracle in the quantum circuit suffices (as here, $M = 1 = N/4$). In other words, one “Grover operator” suffices.
- (b) If P is any projector we have $(I - 2P)(I - 2P) = I - 4P + 4P^2 = I - 4P + 4P = I$. For the given U matrix this implies that $UU^\dagger = U^\dagger U = I$.

The entry $|00\rangle$ is mapped to

$$\begin{aligned} |00\rangle &\rightarrow |11\rangle \rightarrow \frac{1}{\sqrt{2}}(|10\rangle - |11\rangle) \rightarrow \frac{1}{\sqrt{2}}(|11\rangle - |10\rangle) \\ &\rightarrow \frac{1}{2}(|10\rangle - |11\rangle - |10\rangle - |11\rangle) = -|11\rangle \rightarrow -|00\rangle \end{aligned}$$

The entry $|10\rangle$ is mapped to

$$\begin{aligned} |10\rangle &\rightarrow |01\rangle \rightarrow \frac{1}{\sqrt{2}}(|00\rangle - |01\rangle) \rightarrow \frac{1}{\sqrt{2}}(|00\rangle - |01\rangle) \\ &\rightarrow \frac{1}{2}(|00\rangle + |01\rangle - |00\rangle + |01\rangle) = |01\rangle \rightarrow |10\rangle \end{aligned}$$

and we check also that $|01\rangle \rightarrow |01\rangle$ et $|11\rangle \rightarrow |11\rangle$.

- (c) *Algorithmic steps:* We assume that $x_0 = 00$ without loss of generality.

1. Initial state $|001\rangle$
2. $H^{\otimes 3}|001\rangle = \frac{1}{(\sqrt{2})^3}(|00\rangle + |01\rangle + |10\rangle + |11\rangle) \otimes (|0\rangle - |1\rangle)$
3. After the oracle

$$\begin{aligned} &\frac{1}{(\sqrt{2})^3} \{ |00\rangle \otimes (|f(00)\rangle - |\overline{f(00)}\rangle) + |01\rangle \otimes (|f(01)\rangle - |\overline{f(01)}\rangle) \\ &\quad + |10\rangle \otimes (|f(10)\rangle - |\overline{f(10)}\rangle) + |11\rangle \otimes (|f(11)\rangle - |\overline{f(11)}\rangle) \} \end{aligned}$$

Because $f(00) = 1$ and $f(01) = f(10) = f(11) = 0$ we find

$$\begin{aligned} &\frac{1}{(\sqrt{2})^3} \{ |00\rangle \otimes (|1\rangle - |0\rangle) + |01\rangle \otimes (|0\rangle - |1\rangle) \\ &\quad + |10\rangle \otimes (|0\rangle - |1\rangle) + |11\rangle \otimes (|0\rangle - |1\rangle) \} \\ &= \frac{1}{(\sqrt{2})^3} \{ -|00\rangle + |01\rangle + |10\rangle + |11\rangle \} \otimes (|0\rangle - |1\rangle) \end{aligned}$$

Note that the solution $|00\rangle$ is marked here with a phase -1 . This is sometimes called the “kickback phase” phenomenon (like in Deutsch-Josza’s algorithm). Now we apply $H^{\otimes 2}$ to the first register and this gives:

$$\begin{aligned} \frac{1}{(\sqrt{2})^5} \{ & -|00\rangle - |01\rangle - |10\rangle - |11\rangle + |00\rangle - |01\rangle + |10\rangle - |11\rangle \\ & + |00\rangle + |01\rangle - |10\rangle - |11\rangle + |00\rangle - |01\rangle - |10\rangle + |11\rangle \} \otimes (|0\rangle - |1\rangle). \end{aligned}$$

We apply the controlled sign change: only $|00\rangle$ changes sign:

$$\begin{aligned} \frac{1}{(\sqrt{2})^5} \{ & +|00\rangle - |01\rangle - |10\rangle - |11\rangle - |00\rangle - |01\rangle + |10\rangle - |11\rangle \\ & - |00\rangle + |01\rangle - |10\rangle - |11\rangle - |00\rangle - |01\rangle - |10\rangle + |11\rangle \} \otimes (|0\rangle - |1\rangle). \end{aligned}$$

Before proceeding, we simplify:

$$\begin{aligned} \frac{1}{(\sqrt{2})^5} \{ & -2|00\rangle - 2|01\rangle - 2|10\rangle - 2|11\rangle \} \otimes (|0\rangle - |1\rangle) \\ & = -\frac{1}{(\sqrt{2})^3} \{ +|00\rangle + |01\rangle + |10\rangle + |11\rangle \} \otimes (|0\rangle - |1\rangle) \\ & = -\frac{1}{\sqrt{2}} \underbrace{(H^{\otimes 2}|00\rangle)}_{\hat{O} \text{ surprise!}} \otimes (|0\rangle - |1\rangle) = -H^{\otimes 3}(|001\rangle). \end{aligned}$$

4. Now we apply the last series of Hadamard gates $H^{\otimes 3}$. Since $H^2 = 1$ we find the final state $-|00\rangle \otimes \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$. The measurement of the first register gives $x_0 = 00$ with probability 1.