

---

Homework 9 graded  
Introduction to Quantum Information Processing

---

**Exercise 1** *Dynamics of 1-qubit density matrix*

In class we showed that the general form of a 1-qubit density matrix is

$$\rho = \frac{1}{2}(I + \vec{a} \cdot \vec{\sigma})$$

where  $\vec{a} = (a_x, a_y, a_z)$  is a vector in the unit three dimensional ball (the Bloch *ball*)  $\|\vec{a}\| \leq 1$  and  $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$  are the three usual Pauli matrices. Consider the dynamics of this mixed state generated by the Hamiltonian of the qubit in a static plus rotating magnetic field in the rotating frame (as seen in class,  $\omega_1 \propto$  the strength of the rotating field and  $\delta = \omega - \omega_0$  the detuning between the Larmor and rotating field frequencies)

$$H = \frac{\hbar\delta}{2}\sigma_z - \frac{\hbar\omega_1}{2}\sigma_x$$

a) Show that the density matrix at time  $t$  is of the form

$$\rho_t = \frac{1}{2}(I + \vec{a}(t) \cdot \vec{\sigma})$$

and compute the vector  $\vec{a}(t)$ . *Hint:* From the definition of the density matrix you can infer that

$$\rho_t = U_t \rho U_t^\dagger$$

with  $U_t$  the evolution operator.

- b) Check that  $\|\vec{a}(t)\| = \|\vec{a}\|$ . So the vector  $\vec{a}(t)$  evolves on a sphere (inside the Bloch ball) of radius given by the initial vector.
- c) Find a simple proof of the last statement without ever computing  $\vec{a}(t)$ .

**Exercise 2** *The difference between a Bell state and a statistical mixture of  $|00\rangle$  and  $|11\rangle$*

We consider a source that distributes to A and B either an EPR pair in the perfect Bell state  $|B_{00}\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ , or distributes a pair of qubits in a statistical mixture of states  $|00\rangle, |11\rangle$  with uniform probabilities  $1/2$ . This exercise illustrates in many ways that the two kind of situations are completely different.

- a) Write down the density matrix  $\rho_{\text{Bell}}$  associated to the Bell state in Dirac notation as well as in matrix array form (in the computational basis).

- b) Write down the density matrix  $\rho_{stat}$  associated to the statistical mixture above in Dirac notation as well as in matrix array form (in the computational basis).
- c) In a Bell/CSHS experiment one measures the observable

$$\mathcal{B} = A \otimes B + A \otimes B' - A' \otimes B + A' \otimes B'$$

What is the theoretical average if the state when the state is  $\rho_{Bell}$ ? (Use results proven in class and no need to reproduce calculations). And now compute the theoretical average if the state is  $\rho_{stat}$ . What are the values of the of these two averages for the optimal CSHS-angles  $\alpha = 0$ ,  $\alpha' = -\frac{\pi}{4}$ ,  $\beta = \frac{\pi}{8}$ ,  $\beta' = -\frac{\pi}{8}$ ?

### Exercise 3 *Density matrix: a decoherence model*

In the following, we will study a model of decoherence of one qubit interacting with the environment. The whole system is defined in the hilbert space  $\mathcal{H} = \mathcal{H}_{\mathcal{E}} \otimes \mathcal{H}_b$  where  $\mathcal{H}_{\mathcal{E}}$  is the Hilbert space describing the possible states of the environment and  $\mathcal{H}_b = \mathbb{C}^2$  is the Hilbert space describing the possible states of the qubit.

Let  $|\phi_0\rangle = \alpha|0\rangle + \beta|1\rangle \in \mathcal{H}_b$  be the initial state of the qubit and  $|\mathcal{E}\rangle \in \mathcal{H}_b$  that of the environment (or sometimes called *heat-bath*). Let  $(|i\rangle)_{i \geq 1} \in \mathcal{H}_{\mathcal{E}}$  be an "infinite" orthonormal basis of the environment  $\mathcal{H}_{\mathcal{E}}$ . We define the evolution operator  $U = \sum_{i=1}^{+\infty} |i\rangle\langle i| \otimes \mathcal{D}(\theta_i)$  for some distinct angles  $\theta_i \in \mathbb{R}$ , and the dephasing operator:  $\mathcal{D}(\theta_i) = |0\rangle\langle 0| + e^{i\theta_i}|1\rangle\langle 1|$ .

If the environment makes a transition from state  $|\mathcal{E}\rangle$  to  $|i\rangle$ , we let  $\mu(\theta_i) = P(|\mathcal{E}\rangle \rightarrow |i\rangle)$  the probability of such a transition. Note that  $\langle i|\mathcal{E}\rangle = e^{i \arg\langle i|\mathcal{E}\rangle} \sqrt{\mu(\theta_i)}$ .

- a) What is the initial global state  $|\psi_0\rangle$  of the whole system?
- b) Show that  $U$  is a unitary operator (describe your steps).
- c) The state of the system evolves (in discrete time steps say) with a power  $n \in \mathbb{N}$  of the operator  $U$  as  $|\psi_n\rangle = U^n |\psi_0\rangle$ . Show that  $\mathcal{D}(\theta_i)^n = \mathcal{D}(n\theta_i)$  and deduce that

$$|\psi_n\rangle = \sum_{i=1}^{+\infty} e^{i \arg\langle i|\mathcal{E}\rangle} \sqrt{\mu(\theta_i)} |i\rangle \otimes (\mathcal{D}(n\theta_i) |\phi_0\rangle)$$

- d) Now let's consider the density matrix of the qubit itself:  $\rho_n = \text{tr}_{\mathcal{H}_{\mathcal{E}}} [|\psi_n\rangle\langle\psi_n|]$ . First, using only the result of question (a), show that we have initially:

$$\rho_0 = \begin{pmatrix} |\alpha|^2 & \alpha\beta^* \\ \alpha^*\beta & |\beta|^2 \end{pmatrix}$$

- e) For any angle  $\theta \in \mathbb{R}$ , show that we have:

$$\mathcal{D}(\theta)\rho_0\mathcal{D}(\theta)^\dagger = \begin{pmatrix} |\alpha|^2 & \alpha\beta^*e^{-i\theta} \\ \alpha^*\beta e^{i\theta} & |\beta|^2 \end{pmatrix}$$

- f)** Now let's consider  $\hat{\theta}$  a random variable taking values  $\theta_i$  in  $\mathbb{R}$  with probability partial  $\mu(\theta_i)$ . Use the result of question (c) and (e) to show that the density matrix of the qubit coincide with the following expression:

$$\rho_n = \begin{pmatrix} |\alpha|^2 & \alpha\beta^*\mathbb{E}[e^{-in\hat{\theta}}] \\ \alpha^*\beta\mathbb{E}[e^{in\hat{\theta}}] & |\beta|^2 \end{pmatrix}$$

- g)** Now say that the values  $\theta_i$  form a quasicontinuum and that  $\mu$  is the PDF of a gaussian distribution of mean 0 and variance  $\sigma^2$ . Show that the density matrix of the qubit evolves as:

$$\rho_n = \begin{pmatrix} |\alpha|^2 & \alpha\beta^*e^{-\frac{1}{2}\sigma^2n^2} \\ \alpha^*\beta e^{-\frac{1}{2}\sigma^2n^2} & |\beta|^2 \end{pmatrix}$$

Calculate  $\rho_\infty = \lim_{n \rightarrow \infty} \rho_n$ .

- h)** (not graded) How does the von Neumann entropy of the qubit evolve from initial time  $n = 0$  to final time  $n \rightarrow +\infty$ ?