

Astrophysics IV Stellar and galactic dynamics

Exercises

Problem 1:

Derive the Jeans equations in spherical coordinates, starting from the collisionless Boltzmann equation. Assume a spherically symmetric system.

Hints:

- for the zeroth moment, you can express the zeroth moment of the Boltzmann equation

$$\frac{\partial \nu}{\partial t} + \sum_i \frac{\partial}{\partial x_i} (\nu \bar{v}_i) = 0, \quad \nu \equiv \int d^3\mathbf{v} f$$

first in vectorial notation, then in spherical coordinates.

- for the first moment, use the collisionless Boltzmann equation in spherical coordinates, and compute the first moment by multiplying with the radial velocity v_r and integrate over all velocity space.
- Finally, find the solution assuming the system is stationary, isotropic, and express the equation using velocity dispersion σ .
- Here are a few properties to keep in mind :

1) $f \rightarrow 0$ when $|v_i| \rightarrow \infty$ 2) $m \int f d^3\mathbf{v} = \rho$ 3) $m \int v_i f d^3\mathbf{v} = \rho \bar{v}_i$
 4) $m \int v_i v_j f d^3\mathbf{v} = \rho \overline{v_i v_j}$ 5) $\overline{v_i v_j} + \sigma_{ij}^2 = \overline{v_i v_j}$

Problem 2:

Using the Jeans equations in spherical coordinates, determine the velocity dispersion analytically (assuming it is isotropic) for a Plummer sphere in equilibrium.

Problem 3:

Derive the Jeans equations in cylindrical coordinates, starting from the collisionless Boltzmann equation. Assume an axi-symmetric system.