



Differential Geometry II - Smooth Manifolds

Winter Term 2025/2026

Lecturer: Dr. N. Tsakanikas

Assistant: L. E. Rösler

Exercise Sheet 9

Exercise 1 (to be submitted by Thursday, 20.11.2024, 16:00):

- (a) Let $\pi: E \rightarrow M$ be a smooth vector bundle of rank k over a smooth manifold M . Suppose that $\Phi: \pi^{-1}(U) \rightarrow U \times \mathbb{R}^k$ and $\Psi: \pi^{-1}(V) \rightarrow V \times \mathbb{R}^k$ are two smooth local trivializations of E with $U \cap V \neq \emptyset$. Show that the transition function $\tau: U \cap V \rightarrow \text{GL}(k, \mathbb{R})$ between Φ and Ψ is smooth.
- (b) Consider the tangent bundle $\pi: TM \rightarrow M$ of a smooth n -manifold M and let $\Phi: \pi^{-1}(U) \rightarrow U \times \mathbb{R}^n$ and $\Psi: \pi^{-1}(V) \rightarrow V \times \mathbb{R}^n$ be the smooth local trivializations of TM associated with two smooth charts (U, φ) and (V, ψ) for M . Determine the transition function $\tau: U \cap V \rightarrow \text{GL}(n, \mathbb{R})$ between Φ and Ψ .
- (c) Consider the tangent bundle $\pi: TS^2 \rightarrow S^2$ of the unit sphere $S^2 \subseteq \mathbb{R}^3$. Compute the transition function associated with the two local trivializations determined by stereographic coordinates.
- (d) Let $\pi: E \rightarrow M$ be a smooth vector bundle of rank k over a smooth manifold M . Suppose that $\{U_\alpha\}_{\alpha \in A}$ is an open cover of M , and that for each $\alpha \in A$ we are given a smooth local trivialization $\Phi_\alpha: \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{R}^k$ of E . For each $\alpha, \beta \in A$ such that $U_\alpha \cap U_\beta \neq \emptyset$, let $\tau_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}(k, \mathbb{R})$ be the transition function between the smooth local trivializations Φ_α and Φ_β . Show that the following identity is satisfied for all $\alpha, \beta, \gamma \in A$:

$$\tau_{\alpha\beta}(p)\tau_{\beta\gamma}(p) = \tau_{\alpha\gamma}(p) \quad \text{for all } p \in U_\alpha \cap U_\beta \cap U_\gamma. \quad (\star)$$

Exercise 2 (Smooth vector bundle construction lemma):

Let M be a smooth manifold and let $\{U_\alpha\}_{\alpha \in A}$ be an open cover of M . Suppose that for each $\alpha, \beta \in A$ we are given a smooth map $\tau_{\alpha\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}(k, \mathbb{R})$ such that (\star) is satisfied for all $\alpha, \beta, \gamma \in A$. Show that there is a smooth vector bundle $E \rightarrow M$ of rank k with smooth local trivializations $\Phi_\alpha: \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{R}^k$ whose transition functions are the given maps $\tau_{\alpha\beta}$.

[Hint: Define an appropriate equivalence relation on $\coprod (U_\alpha \times \mathbb{R}^k)$ and use the vector bundle chart lemma.]

Exercise 3:

- (a) Show that the zero section of every smooth vector bundle is smooth.

[Hint: Consider $\Phi \circ \zeta$, where Φ is a smooth local trivialization.]

- (b) Let $\pi: E \rightarrow M$ be a smooth vector bundle. Show that if $f, g \in C^\infty(M)$ and if $\sigma, \tau \in \Gamma(E)$, then $f\sigma + g\tau \in \Gamma(E)$.

[Hint: Consider $\Phi \circ (f\sigma + g\tau)$, where Φ is a smooth local trivialization of E .]

- (c) Let $E := M \times \mathbb{R}^k$ be a product bundle over a topological manifold M . Show that there is a natural one-to-one correspondence between continuous sections of E and continuous functions from M to \mathbb{R}^k .

Moreover, if M is a smooth manifold, show that this is a one-to-one correspondence between smooth sections of E and smooth functions from M to \mathbb{R}^k . Deduce that there is a natural identification between the space $C^\infty(M)$ and the space of smooth sections of the trivial line bundle $M \times \mathbb{R} \rightarrow M$.

Exercise 4 (*Completion of smooth local frames for smooth vector bundles*):

Let $\pi: E \rightarrow M$ be a smooth vector bundle of rank k over a smooth manifold M . Prove the following assertions:

- (a) If $(\sigma_1, \dots, \sigma_m)$ is a linearly independent m -tuple of smooth local sections of E over an open subset $U \subseteq M$, where $1 \leq m < k$, then for each $p \in U$ there exist smooth sections $\sigma_{m+1}, \dots, \sigma_k$ of E defined on some neighborhood V of p such that $(\sigma_1, \dots, \sigma_k)$ is a smooth local frame for E over $U \cap V$.
- (b) If (v_1, \dots, v_m) is a linearly independent m -tuple of elements of the fiber E_p for some $p \in M$, where $1 \leq m < k$, then there exists a smooth local frame $(\sigma_1, \dots, \sigma_k)$ for E over some neighborhood of p such that $\sigma_i(p) = v_i$ for every $1 \leq i \leq m$.
- (c) If $A \subseteq M$ is a closed subset and if (τ_1, \dots, τ_k) is a linearly independent k -tuple of sections of $E|_A$ which are smooth in the sense described in *Lemma 6.11*, then there exists a smooth local frame $(\sigma_1, \dots, \sigma_k)$ for E over some neighborhood of A such that $\sigma_i|_A = \tau_i$ for every $1 \leq i \leq k$.

[Hint: Use *Lemma 6.11*.]

Exercise 5 (*Correspondence between smooth local frames and smooth local trivializations*): Let $\pi: E \rightarrow M$ be a smooth vector bundle of rank k over a smooth n -manifold M .

- (a) Given a smooth local trivialization $\Phi: \pi^{-1}(U) \rightarrow U \times \mathbb{R}^k$ of E over U , construct a smooth local frame (σ_i) for E over U . (We say that the smooth local frame (σ_i) is associated with the smooth local trivialization Φ .)
- (b) Show that every smooth local frame (σ_i) for E is associated with a smooth local trivialization Φ of E .

[Hint: Define the inverse of Φ using (σ_i) and show that it is a bijective local diffeomorphism to conclude.]

- (c) Deduce that E is smoothly trivial if and only if it admits a smooth global frame. Interpret this result in case that E is a smooth *line bundle*, i.e., when $k = 1$.
- (d) Let (U, φ) be a smooth coordinate chart for M with coordinate functions (x^i) and assume that there exists a smooth local frame (σ_i) for E over U . Consider the map

$$\tilde{\varphi}: \pi^{-1}(U) \rightarrow \varphi(U) \times \mathbb{R}^k, \quad v^i \sigma_i(p) \mapsto (x^1(p), \dots, x^n(p), v^1, \dots, v^k).$$

Show that $(\pi^{-1}(U), \tilde{\varphi})$ is a smooth coordinate chart for E .

Exercise 6 (*Uniqueness of the smooth structure on TM*):

Let M be a smooth n -manifold. Show that the topology and smooth structure on the tangent bundle TM constructed in *Proposition 3.12* are the unique ones with respect to which $\pi: TM \rightarrow M$ is a smooth vector bundle with the given vector space structure on the fibers, and such that all coordinate vector fields are smooth local sections.

[Hint: Use *Exercise 5(d)*.]